



10-30-00

Box SEQ A

Patent Application Transmittal
(only for new nonprovisional applications
under 37 C.F.R. 1.53(b))
Correspondence Address:
FROMMER LAWRENCE & HAUG LLP
745 Fifth Avenue
New York, New York 10151
Tel. (212) 588-0800
Fax: (212) 588-0500



Date: October 27, 2000
Attorney Ref.: 514412-2020.1

TO: ASSISTANT COMMISSIONER FOR PATENTS
Box PATENT APPLICATION
Washington, D.C. 20231

Sir:

With reference to the filing in the United States Patent and Trademark Office of an application for patent in the name of:

WESTON ET AL.

Entitled: MALE-STERILE *BRASSICA* PLANTS AND METHODS FOR PRODUCING SAME

- This is an application of a small entity under 37 CFR 1.9(f) and the amounts shown in parentheses below have been employed in calculating the fee.
- Small Entity Verified Statement is enclosed (unsigned).
- Small Entity Verified Statement filed in prior application, status still proper and desired

The following are enclosed:

- X Specification (41 pages) (One Page Abstract (Page 48)).
- X 5 Sheet(s) of Drawings
- X 22 Claims (including 8 independent and 12 dependent claims) (Pages 42-47 of application)
- X 7 pages Sequence Listing
- This application contains multiple dependent claim(s).

09698903-102700

X Our check totaling \$1,136.00, calculated on the basis of the attached Continuation-In-Part Application of U.S. patent application Serial No. 09/430,437 filed October 29, 1999 as follows:

Basic Fee
\$710.00 (355.00).....710.00

Total Number of Claims in excess
of 20 at \$18.00 (9.00) each36.00

Number of Independent Claims in
excess of 3 at \$78.00 (39.00) each.....390.00

Multiple Dependent Claim Fee
at \$270.00 (135.00).....0.00

Total Filing Fee.....1,136.00

Assignment Recording Fee \$40.000.00

X Oath or Declaration and Power of Attorney

_____ New _____ signed X unsigned

_____ Copy from prior application (37 C.F.R. 1.63(d))

_____ Certified copy of the following application to substantiate the claim for priority made in the Declaration:

Please charge any additional fees required for the filing of this application or credit any overpayment to Deposit Account No. 50-0320.

Respectfully submitted,

FROMMER LAWRENCE & HAUG LLP

By: Susan Lehnhardt
Susan K. Lehnhardt
Reg. No. 33,943
(212) 588-0800

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE
APPLICATION FOR LETTERS PATENT

JC920 U.S. PTO
09/698903
10/27/00

TITLE : MALE-STERILE *BRASSICA* PLANTS AND METHODS FOR
PRODUCING SAME

INVENTOR : WESTON ET AL.

- 41 Pages specification
- 6 Claims (pages 42 to 47, including 8 independent claims)
- 5 Figs. (5 sheets) (Figs 1, 2, 3, 4, 5)
- 1 page(s) Abstract (p. 48)
- 7 pages of Sequence Listing

EXPRESS MAIL

Mailing Label Number: EL588272530US

Date of Deposit: October 27, 2000

I hereby certify that this paper or fee is being deposited with the United States Postal Service "Express Mail Post Office to Addressee" Service under 37 CFR 1.10 on the date indicated above and is addressed to the Commissioner of Patents and Trademarks, Washington, DC 20231.

Charles Jackson
(Typed or printed name of person mailing paper or fee)

Charles Jackson
(Signature of person mailing paper or fee)

William F. Lawrence
Reg. No. 28,029
Susan K. Lehnhardt
Reg. No. 33,943
FROMMER LAWRENCE & HAUG LLP
745 Fifth Avenue
New York, New York 10151
Phone: 212 588 0800
Fax: 212 588-0500

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Applicant(s) : WESTON et al.
U.S. Serial No. : N/A
Filing Date : N/A
For : MALE-STERILE *BRASSICA* PLANTS AND
METHODS FOR PRODUCING SAME

745 Fifth Avenue
New York, New York 10151

EXPRESS MAIL

Mailing Label Number: EL588272530US

Date of Deposit: October 27, 2000

I hereby certify that this paper or fee is being deposited with the United States Postal Service "Express Mail Post Office to Addressee" Service under 37 CFR 1.10 on the date indicated above and is addressed to the Assistant Commissioner for Patents, Washington, D.C. 20231.

Charles Jackson
(Typed or printed name of person mailing paper or fee)

Charles Jackson
(Signature of person mailing paper or fee)

**STATEMENT TO SUPPORT FILING AND
SUBMISSION IN ACCORDANCE WITH 37 C.F.R. §§ 1.821-1.825**

Box Sequence
Assistant Commissioner for Patents
Washington, D.C. 20231

Dear Sir:

The undersigned hereby states that the content of the printed Sequence Listing for the above-referenced application filed, and the computer readable copy, submitted in accordance with 37 C.F.R. §§ 1.821(c) and (e), are the same.

In the unlikely event that the Patent Office determines that an extension and/or other relief is required as a result of this statement, applicants petition for any required relief

Susan K. Lehnhardt
Susan K. Lehnhardt
Reg. No. 33,943
(212) 588-0800

INVENTORS

Full name of first inventor: WESTON, Brigitte
 Residence: Fatimastraat 13,
B-9880 Aalter
Belgium
 Post Office: same as residence
 Citizenship: Canadian

Full name of second inventor: De BEUCKLELEER, Marc
 Residence: Heckers 36
B-9052 Zwijnaarde
Belgium
 Post Office: same as residence
 Citizenship: Belgian

	1970	1971	1972	1973	1974	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024	2025	2026	2027	2028	2029	2030	2031	2032	2033	2034	2035	2036	2037	2038	2039	2040	2041	2042	2043	2044	2045	2046	2047	2048	2049	2050	2051	2052	2053	2054	2055	2056	2057	2058	2059	2060	2061	2062	2063	2064	2065	2066	2067	2068	2069	2070	2071	2072	2073	2074	2075	2076	2077	2078	2079	2080	2081	2082	2083	2084	2085	2086	2087	2088	2089	2090	2091	2092	2093	2094	2095	2096	2097	2098	2099	2100	2101	2102	2103	2104	2105	2106	2107	2108	2109	2110	2111	2112	2113	2114	2115	2116	2117	2118	2119	2120	2121	2122	2123	2124	2125	2126	2127	2128	2129	2130	2131	2132	2133	2134	2135	2136	2137	2138	2139	2140	2141	2142	2143	2144	2145	2146	2147	2148	2149	2150	2151	2152	2153	2154	2155	2156	2157	2158	2159	2160	2161	2162	2163	2164	2165	2166	2167	2168	2169	2170	2171	2172	2173	2174	2175	2176	2177	2178	2179	2180	2181	2182	2183	2184	2185	2186	2187	2188	2189	2190	2191	2192	2193	2194	2195	2196	2197	2198	2199	2200	2201	2202	2203	2204	2205	2206	2207	2208	2209	2210	2211	2212	2213	2214	2215	2216	2217	2218	2219	2220	2221	2222	2223	2224	2225	2226	2227	2228	2229	2230	2231	2232	2233	2234	2235	2236	2237	2238	2239	2240	2241	2242	2243	2244	2245	2246	2247	2248	2249	2250	2251	2252	2253	2254	2255	2256	2257	2258	2259	2260	2261	2262	2263	2264	2265	2266	2267	2268	2269	2270	2271	2272	2273	2274	2275	2276	2277	2278	2279	2280	2281	2282	2283	2284	2285	2286	2287	2288	2289	2290	2291	2292	2293	2294	2295	2296	2297	2298	2299	2300	2301	2302	2303	2304	2305	2306	2307	2308	2309	2310	2311	2312	2313	2314	2315	2316	2317	2318	2319	2320	2321	2322	2323	2324	2325	2326	2327	2328	2329	2330	2331	2332	2333	2334	2335	2336	2337	2338	2339	2340	2341	2342	2343	2344	2345	2346	2347	2348	2349	2350	2351	2352	2353	2354	2355	2356	2357	2358	2359	2360	2361	2362	2363	2364	2365	2366	2367	2368	2369	2370	2371	2372	2373	2374	2375	2376	2377	2378	2379	2380	2381	2382	2383	2384	2385	2386	2387	2388	2389	2390	2391	2392	2393	2394	2395	2396	2397	2398	2399	2400	2401	2402	2403	2404	2405	2406	2407	2408	2409	2410	2411	2412	2413	2414	2415	2416	2417	2418	2419	2420	2421	2422	2
--	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	---

TITLE OF THE INVENTION

MALE-STERILE *BRASSICA* PLANTS AND METHODS FOR PRODUCING SAME

CROSS-REFERENCE TO RELATED APPLICATIONS

This application is a continuation-in-part of US patent application Serial No.

5 09/430,437, filed October 29, 1999, pending.

STATEMENT REGARDING FEDERALLY SPONSORED RESEARCH OR DEVELOPMENT

Not applicable.

FIELD OF THE INVENTION

10 This invention pertains to transgenic *Brassica* plants, plant material and seeds, characterized by harboring a specific transformation event, particularly by the presence of a male-sterility gene, at a specific location in the *Brassica* genome. The *Brassica* plants of the invention combine the male-sterility phenotype with optimal agronomic performance, genetic stability and adaptability to different genetic backgrounds.

15 All documents cited herein are hereby incorporated herein by reference.

BACKGROUND OF THE INVENTION

20 The phenotypic expression of a transgene in a plant is determined both by the structure of the gene itself and by its location in the plant genome. At the same time the presence of the transgene (in a foreign DNA) at different locations in the genome will influence the overall phenotype of the plant in different ways. The agronomically or industrially successful introduction of a commercially interesting trait in a plant by genetic manipulation can be a lengthy procedure dependent on different factors. The actual transformation and regeneration of genetically transformed plants are only the first

in a series of selection steps that include extensive genetic characterization, breeding, and evaluation in field trials.

The term “rapeseed” covers every seed of the *Brassica* species. *Brassica* is cultivated from China and India to Finland and Canada as one of the most valuable oil crops. Most *Brassica* types belong to the family of *Cruciferae*. They originated as a diploid species having aneuploid chromosome numbers ranging from 7 (*Brassica fruticulosa*) to 12 (*Sinapsis alba*).

The most extensively grown *Brassica* species in Canada is known as turnip rape, or *Brassica campestris* (aa, n=10). *Brassica oleracea* (cc, n=9) has diversified in recent evolutionary history into at least six major horticultural types, including broccoli, cauliflower and cabbage. *Brassica nigra* (bb, n= 8) or black mustard is a less important crop commercially and is mainly known for its seeds from which mustard was originally made. From these basic types, amphiploid hybrids have been derived in more recent evolutionary stages by intercrossing. The most important of these are *Brassica napus* (aacc), of which the winter types provide the highest rapeseed yields in northern Europe and *Brassica juncea* (aabb) or brown mustard that is one of the major oil crops of the Indian sub-continent.

Though intercrossing between different *Brassica* species (particularly those with compatible genomes) is possible and often done for breeding purposes, not all traits (or genes) will be able to be transferred from one species to the other or, when transferred to a different species, will not retain identical characteristics (or expression patterns). Thus, a genetic locus conferring optimal expression of a natural or chimeric gene in one *Brassica* species, will not necessarily have the same effect in another.

Brassica species are bisexual and typically 60-70% self-pollinated. The production of hybrids and introduction of genetic variation as a basis for selection was traditionally dependent on the adaptation of natural occurring phenomena such as self-incompatibility and cytoplasmic male-sterility. Artificial pollination control methods
5 such as manual emasculation or the use of gametocides are not widely applied in *Brassica* breeding due to their limited practicability and high cost respectively.

Transgenic methods have been developed for the production of male or female-sterile plants, which provide interesting alternatives to the traditional techniques.

EP 0,344,029 describes a system for obtaining nuclear male-sterility where plants
10 are transformed with a male-sterility gene, comprising for example a DNA encoding a barnase molecule under the control of a tapetum specific promoter TA29, which when incorporated into a plant ensures selective destruction of tapetum cells. Transformation of tobacco and oilseed rape plants with such a gene resulted in plants in which pollen formation was completely prevented. Mariani et al. (1990) Nature 347:737-741.

15 Cytochemical and histochemical analysis of anther development of *B. napus* plants comprising the chimeric PTA29:barnase gene is described by De Block and De Brouwer ((1993) Planta 189:218-225).

To restore fertility in the progeny of a male-sterile plant, a system was developed where the male-sterile plant is crossed with a transgenic plant carrying a fertility-restorer
20 gene, which when expressed is capable of inhibiting or preventing the activity of the male-sterility gene. US Patent Nos. 5,689,041; and 5,792,929; and De Block and De Brouwer (1993).

The use of coregulating genes in the production of male-sterile plants to increase the frequency of transformants having good agronomic performance is described in WO 96/26283. Typically, when the sterility DNA encodes a barnase, the coregulating DNA will encode a barstar.

5 Successful genetic transformation of *Brassica* species has been obtained by a number of methods including *Agrobacterium* infection (as described, for example in EP 0,116,718 and EP 0,270,882), microprojectile bombardment (as described, for example by Chen et al. (1994) Theor. Appl. Genet. 88:187-192) and direct DNA uptake into protoplasts (as described, for example by De Block et al. (1989) Plant Physiol. 914:694-701; and Poulsen (1996) Plant Breeding 115:209-225).

However, the foregoing documents fail to teach or suggest the present invention.

SUMMARY OF THE INVENTION

The invention relates to a transgenic *Brassica* plant, the genomic DNA of which is characterized by one or both of the following characteristics:

a) the genomic DNA is capable of yielding at least two, preferably at least three, more preferably at least four, most preferably five of the restriction fragments or sets of restriction fragments selected from the group of:

i) One set of NcoI fragments, one with a length of between 5077 and 14057 bp, preferably of about 6000 bp, and one with a length of between 2450 and 2838 bp, preferably of about 2500 bp;

ii) one set of EcoRV fragments wherein one has a length of between 5077 and 14057 bp, preferably of about 5500 bp and one with a length of between 4507 and 5077 bp, preferably of about 4800 bp;

iii) one set of MunI fragments, one with a length of between 5077 and 14057 bp, preferably with a length of about 5700 bp, and one with a length of between 2838 and 4799 bp, preferably of about 4500 bp;

iv) one HindIII fragment, with a length of between 2838 and 4507 bp, preferably with a length of about 3938 bp,

v) one EcoRI fragment, with a length of between 1989 and 2450 bp, preferably of about 2262 bp;

wherein each of the restriction fragments is capable of hybridizing under standard stringency conditions, with the +/- 2000 bp fragment obtainable by PCR amplification of a fragment of SEQ ID NO:1, using the probes having SEQ ID NO:2 and SEQ ID NO:3 and/or

b) the genomic DNA can be used to amplify a DNA fragment of between 160 and 200 bp, preferably of about 183 bp, using a polymerase chain reaction with two primers having the nucleotide sequence of SEQ ID NO:11 and SEQ ID NO:12 respectively.

The present invention further relates to a transgenic *Brassica* plant, or seed, cells or tissues thereof, the genomic DNA of which is characterized in that it is capable of yielding at least two, preferably at least three, for instance at least four, more preferably five of the sets of restriction fragments selected from the group described under a) above comprising the sets of restriction fragments described under a) i), ii), iii), iv), and v) above, whereby the selection can include any combination of i), ii), iii), iv), and v) described under a) above.

The present invention further relates to a transgenic *Brassica* plant, or seed, cells, tissues or progeny thereof, the genomic DNA of which is characterized by both the characteristics described under a) and b) above.

The present invention further relates to a transgenic male-sterile *Brassica* plant, the genomic DNA of which is characterized by one, preferably by both the characteristics described under a) and b) above.

The invention also relates to the seed deposited at the ATCC under number PTA-850 or PTA-2485, which will grow into a male-sterile, herbicide resistant plant. The seed of ATCC deposit number PTA-850 or PTA-2485 comprises about 50% seed comprising the elite event of the invention, which will grow into male-sterile, PPT tolerant plants. The seed can be sown and the growing plants can be treated with PPT or LibertyTM as described herein to obtain 100% male-sterile, PPT tolerant plants, comprising the elite event of the invention. The invention further relates to cells, tissues, progeny, and descendants from a plant comprising the elite event of the invention grown from the seed deposited at the ATCC having accession number PTA-850 or PTA-2485. The invention further relates to plants obtainable by propagation of and/or breeding with a *Brassica* plant comprising the elite event of the invention grown from the seed deposited at the ATCC having accession number PTA-850 or PTA-2485.

The invention further relates to plants, seeds, cells or tissues comprising a foreign DNA sequence, preferably a male-sterility gene as described herein, integrated into the chromosomal DNA in a region which comprises the plant DNA sequence of SEQ ID NO:8 and/or SEQ ID NO:10, or a sequence which has at least 85% sequence identity to a sequence comprising the plant DNA sequence of SEQ ID NO:8 and/or SEQ ID NO:10.

The invention further provides a process for producing a transgenic cell of a *Brassica* plant, which comprises inserting a recombinant DNA molecule into a region of the chromosomal DNA of a *Brassica* cell which comprises the plant DNA sequence of SEQ ID NO:8 and/or SEQ ID NO:10, or a sequence which has at least 85% sequence identity with a sequence comprising the plant DNA sequence of SEQ ID NO:8 and/or SEQ ID NO:10, and, optionally, regenerating a *Brassica* plant from the transformed *Brassica* cell.

The invention further relates to a method for identifying a transgenic plant, or cells or tissues thereof, which method comprises establishing one or both of the following characteristics of the genomic DNA of the transgenic plant, or its cells or tissues:

a) the genomic DNA is capable of yielding at least two, preferably at least three, more preferably at least four, most preferably five of the restriction fragments or sets of restriction fragments selected from the group of:

i) One set of NcoI fragments, one with a length of between 5077 and 14057 bp, preferably of about 6000 bp, and one with a length of between 2450 and 2838 bp, preferably of about 2500 bp;

ii) one set of EcoRV fragments wherein one has a length of between 5077 and 14057 bp, preferably of about 5500 bp and one with a length of between 4507 and 5077 bp, preferably of about 4800 bp;

iii) one set of MunI fragments, one with a length of between 5077 and 14057 bp, preferably with a length of about 5700 bp, and one with a length of between 2838 and 4799 bp, preferably of about 4500 bp;

iv) one HindIII fragment, with a length of between 2838 and 4507 bp,
preferably with a length of about 3938 bp,

v) one EcoRI fragment, with a length of between 1989 and 2450 bp,
preferably of about 2262 bp;

5 wherein each of the restriction fragments is capable of hybridizing under standard stringency conditions, with the +/- 2000 bp fragment obtainable by PCR amplification of a fragment of SEQ ID NO:1, using the probes having SEQ ID NO:2 and SEQ ID NO:3 and/or

b) the genomic DNA can be used to amplify a DNA fragment of between
10 160 and 200 bp, preferably of about 183 bp, using a polymerase chain reaction with two primers having the nucleotide sequence of SEQ ID NO:11 and SEQ ID NO:12 respectively.

The invention further relates to a kit for identifying the transgenic plants comprising the elite event of the present invention, which kit comprises at least two PCR
15 probes, one of which recognizes a sequence within the T-DNA of the elite event, the other recognizing a sequence within the 5' or 3' border flanking region of the elite event of the invention, preferably the PCR primers having the nucleotide sequence of SEQ ID NO:11 and SEQ ID NO:12, respectively for use in the PCR identification protocol.

BRIEF DESCRIPTION OF THE DRAWINGS

20 The following detailed description, given by way of example, but not intended to limit the invention to specific embodiments described, may be understood in conjunction with the accompanying Figures, incorporated herein by reference, in which:

Figure 1. Plasmid map of pDE110. The plasmid pDE110 comprises the coding sequence of the bialaphos resistance gene (bar) of *Streptomyces hygroscopicus*, (Thompson et al. (1987 EMBO J. 6:2519-2523) under control of the promoter of the 35S gene from Cauliflower Mosaic Virus. Odell et al. (1985) Nature 313:810-812. The BglIII-NcoI restriction fragment provides the bar probe.

Figure 2. Plasmid map of pCO48. The plasmid pCO48 comprises the coding sequence of the barnase gene from *Bacillus amyloliquefaciens* (Hartley (1988) J. Mol. Biol. 202:913-915), under control of the promoter region of the anther specific gene TA29 from *Nicotiana tabacum*. Seurinck et al. (1990) Nucl. Acids Res. 18:3403. The EcoRV-NsiI restriction fragment provides the TA29 probe.

Figure 3. Restriction map obtained after digestion of MS-B2 genomic DNA. Loading sequence of the gel analyzed by Southern blot: lane 1, MS-B2 DNA digested with NcoI, lane 2, MS-B2 DNA digested with EcoRV, lane 3, MS-B2 DNA digested with MunI, lane 4, MS-B2 DNA digested with HindIII, lane 5, MS-B2 DNA digested with EcoRI, lane 6, non-transgenic *B. napus* DNA digested with EcoRI, lane 7, non-transgenic *B. napus* DNA digested with EcoRI + control plasmid pTCO113 DNA digested with EcoRI, lane 8, Molecular weight marker (λ - PstI).

Figure 4. PCR analysis of other events and elite event MS-B2 using the MS-B2 PCR identification protocol. Loading sequence of the gel: lane 1, DNA sample from a *Brassica* plant comprising the transgenic event MS-B2, lane 2-5, DNA samples from *Brassica* plants comprising other transgenic events, lane 6, DNA from wild-type *Brassica*, lane 7, negative control (water), lane 8, molecular weight marker (100bp ladder).

DETAILED DESCRIPTION OF THE INVENTION

The term “gene” as used herein refers to any DNA sequence comprising several operably linked DNA fragments such as a promoter and a 5’ untranslated region (the 5’UTR), which together form the promoter region, a coding region (which may or may not code for a protein), and an untranslated 3’ region (3’UTR) comprising a polyadenylation site. Typically in plant cells, the 5’UTR, the coding region and the 3’UTR are transcribed into an RNA that, in the case of a protein-encoding gene, is translated into the protein. A gene may include additional DNA fragments such as, for example, introns. As used herein, a genetic locus is the position of a given gene in the genome of a plant.

The term “chimeric” when referring to a gene or DNA sequence is used to refer to the fact that the gene or DNA sequence comprises at least two functionally relevant DNA fragments (such as promoter, 5’UTR, coding region, 3’UTR, intron) that are not naturally associated with each other and originate, for example, from different sources. “Foreign” referring to a gene or DNA sequence with respect to a plant species is used to indicate that the gene or DNA sequence is not naturally found in that plant species, or is not naturally found in that genetic locus in that plant species. The term “foreign DNA” will be used herein to refer to a DNA sequence as it has incorporated into the genome of a plant as a result of transformation. The “transforming DNA” as used herein refers to a recombinant DNA molecule used for transformation. The transforming DNA usually comprises at least one “gene of interest” (e.g. a chimeric gene) which is capable of conferring one or more specific characteristics to the transformed plant. The term “recombinant DNA molecule” is used to exemplify and thus can include an isolated

nucleic acid molecule which can be DNA and which can be obtained through recombinant or other procedures.

As used herein the term “transgene” refers to a gene of interest as incorporated in the genome of a plant. A “transgenic plant” refers to a plant comprising at least one
5 transgene in the genome of all of its cells.

The foreign DNA present in the plants of the present invention preferably comprises two genes of interest, more specifically, a male-sterility gene and a coregulating gene.

A “male-sterility gene” as used herein refers to a gene that upon expression in a
10 plant renders the plant incapable of producing fertile pollen. An example of a male-sterility gene is a gene comprising a DNA sequence encoding barnase, under the control of a promoter directing expression in tapetum cells. More specifically, according to the present invention, a preferred embodiment of the male-sterility gene comprises a DNA
15 encoding barnase under control of the promoter of the TA29 gene of *N. tabacum*, also referred to as “TA29-barnase” herein. The TA29 promoter has a “tapetum selective” expression pattern in *Brassica*. De Block and Debrouwer (1993).

A “coregulating gene” as used herein refers to a gene, which when introduced into plant cells together with another chimeric gene, increases the frequency of transformants having good agronomical performance. An example of a coregulating gene for use
20 together with a male-sterility gene encoding barnase, is a gene comprising a DNA sequence encoding barstar, under the control of a promoter capable of directing expression in non-stamen cells or which directs weak expression in stamen cells. More specifically, in the elite event of the present invention the coregulating gene comprises a

DNA encoding barstar under control of the nopaline synthase gene from the T-DNA of *Agrobacterium tumefaciens* (Depicker et al. (1984) J. Mol. Appl. Genet. 1:3403), also referred to as “PNOS-barstar” herein.

A “fertility restorer gene” as used herein refers to a gene that upon expression in a plant comprising a male-sterility gene, is capable of preventing expression of the male-sterility gene, restoring fertility in the plant.

The incorporation of a recombinant DNA molecule in the plant genome typically results from transformation of a cell or tissue (or from another genetic manipulation).

The particular site of incorporation is either due to chance or is at a predetermined location (if a process of targeted integration is used).

The foreign DNA can be characterized by the location and the configuration at the site of incorporation of the recombinant DNA molecule in the plant genome. The site in the plant genome where a recombinant DNA has been inserted is also referred to as the “insertion site” or “target site”. Insertion of the recombinant DNA into the plant genome can be associated with a deletion of plant DNA, referred to as “target site deletion”. A “flanking region” or “flanking sequence” as used herein refers to a sequence of at least 20 bp, preferably at least 50 bp, and up to 5000 bp of the plant genome which is located either immediately upstream of and contiguous with or immediately downstream of and contiguous with the foreign DNA. Transformation procedures leading to random integration of the foreign DNA will result in transformants with different flanking regions, which are characteristic and unique for each transformant. When the recombinant DNA is introduced into a plant through traditional crossing, its insertion site in the plant genome, or its flanking regions will generally not be changed.

An “insertion region” as used herein refers to the region corresponding to the region of at least 40 bp, preferably at least 100 bp, and up to more than 10000 bp, encompassed by the sequence which comprises the upstream and/or the downstream flanking region of a foreign DNA in the (untransformed) plant genome (and possibly including the insertion site and possible target site deletion). Taking into consideration minor differences due to mutations within a species, an insertion region will retain at least 85%, preferably 90%, more preferably 95%, and most preferably 100% sequence identity with the sequence comprising the upstream and downstream flanking regions of the foreign DNA in a given plant of that species.

Expression of a gene of interest refers to the fact that the gene confers on the plant one or more phenotypic traits (e.g. herbicide tolerance) that were intended to be conferred by the introduction of the recombinant DNA molecule – the transforming DNA - used during transformation (on the basis of the structure and function of part or all of the gene(s) of interest).

An “event” is defined as a (artificial) genetic locus that, as a result of genetic manipulation, carries a foreign DNA comprising at least one copy of the gene(s) of interest. The typical allelic states of an event are the presence or absence of the foreign DNA. As used herein an “MS” event will refer to an event carrying a transgene comprising “TA29-barnase” and “PNOS-barstar”. An event is characterized phenotypically by the expression of the transgenes. At the genetic level, an event is part of the genetic makeup of a plant. At the molecular level, an event is characterized by the restriction map (e.g. as determined by Southern blotting) and/or by the upstream and/or downstream flanking sequences of the foreign DNA, and/or the molecular configuration

of the foreign DNA comprising the transgenes. Usually transformation of a plant with a transforming DNA leads to a multitude of events, each of which is unique.

An “elite event”, as used herein, is an event which is selected from a group of events, obtained by transformation with the same transforming DNA or by back-crossing with plants obtained by such transformation, based on the expression and stability of the transgenes and its compatibility with optimal agronomic characteristics of the plant comprising it. Thus the criteria for elite event selection are one or more, preferably two or more, advantageously all of the following:

a) That the presence of the foreign DNA does not compromise other desired characteristics of the plant, such as those relating to agronomic performance or commercial value;

b) That the event is characterized by a well defined molecular configuration which is stably inherited and for which appropriate diagnostic tools for identity control can be developed; and

c) That the gene(s) of interest show(s) a correct, appropriate and stable spatial and temporal phenotypic expression, both in heterozygous (or hemizygous) and homozygous condition of the event, at a commercially acceptable level in a range of environmental conditions in which the plants carrying the event are likely to be exposed in normal agronomic use.

It is preferred that the foreign DNA is associated with a position in the plant genome that allows introgression into desired commercial genetic backgrounds.

The status of an event as an elite event is confirmed by introgression of the elite event in different relevant genetic backgrounds and observing compliance with one, two or all of the criteria e.g. a), b) and c) above.

Additionally, for the transforming DNA comprising a male-sterility gene and a coregulating gene described herein, selection of the elite event will also be determined on the compatibility between this event and selected plants comprising a fertility restorer gene. More specifically, it will be ensured that the progeny, resulting from a cross between a plant carrying the male-sterility event as described herein and a plant homozygous for a fertility restorer gene, that this progeny in which at least the fertility restorer event is present, has the following characteristics:

- a) adequate phenotypic expression of the fertility restored phenotype, i.e. male-fertility; and
- b) phenotypic expression occurs at a commercially acceptable level in a range of environmental conditions in which plants carrying the two events are likely to be exposed in normal agronomic use.

An “elite event” thus refers to a genetic locus comprising a foreign DNA, which answers to the above-described criteria. A plant, plant material or progeny such as seeds can comprise one or more elite events in its genome.

The “diagnostic tools” developed to identify an elite event or the plant or plant material comprising an elite event, are based on the specific genomic characteristics of the elite event, such as, a specific restriction map of the genomic region comprising the foreign DNA and/or the sequence of the flanking region(s) of the foreign DNA. A “restriction map” as used herein refers to a set of Southern blot patterns obtained after

cleaving plant genomic DNA with a particular restriction enzyme, or set of restriction enzymes and hybridization with a probe sharing sequence similarity with the foreign DNA under standard stringency conditions. Standard stringency conditions as used herein refers to the conditions for hybridization described herein or to the conventional hybridizing conditions as described by Sambrook et al. (1989) (Molecular Cloning: A Laboratory Manual, Second Edition, Cold Spring Harbor Laboratory Press, NY) which for instance can comprise the following steps: 1) immobilizing plant genomic DNA fragments on a filter, 2) prehybridizing the filter for 1 to 2 hours at 42°C in 50% formamide, 5 X SSPE, 2 X Denhardt's reagent and 0.1% SDS, or for 1 to 2 hours at 68°C in 6 X SSC, 2 X Denhardt's reagent and 0.1% SDS, 3) adding the hybridization probe which has been labeled, 4) incubating for 16 to 24 hours, 5) washing the filter for 20 min. at room temperature in 1X SSC, 0.1 %SDS, 6) washing the filter three times for 20 min. each at 68°C in 0.2 X SSC, 0.1 %SDS, and 7) exposing the filter for 24 to 48 hours to X-ray film at -70°C with an intensifying screen.

Due to the (endogenous) restriction sites present in a plant genome prior to incorporation of the foreign DNA, insertion of a foreign DNA will alter the specific restriction map of that genome. Thus, a particular transformant or progeny derived thereof can be identified by one or more specific restriction patterns. The conditions for determining the restriction map of an event are laid out in a "restriction map identification protocol". Alternatively, plants or plant material comprising an elite event can be identified by testing according to a PCR identification protocol. This is a PCR using primers that specifically recognize the elite event. Essentially, a set of primers is developed which recognizes a) a sequence within the 3' or 5' flanking sequence of the

elite event and b) a sequence within the foreign DNA, which primers amplify a fragment (integration fragment) preferably of between 100 and 300 nucleotides. Preferably, a control is included of a set of primers that amplifies a fragment within a housekeeping gene of the plant species (preferably a fragment which is larger than the amplified integration fragment). The optimal conditions for the PCR, including the sequence of the specific primers are specified in a PCR identification protocol.

The present invention relates to the development of an elite event in *Brassica*, MS-B2, to the plants comprising this event, the progeny obtained from these plants and to the plant cells, or plant material derived from this event. Plants comprising elite event MS-B2 were obtained through transformation with pTCO113 as described in Example 1.

The recombinant DNAs used for the generation of the plants leading to the selection of elite event MS-B2 additionally comprise a DNA sequence encoding the enzyme phosphinothricin acetyl transferase and the 35S promoter of Cauliflower Mosaic Virus (termed “35S-bar”), where the sequence encoding phosphinothricin acetyl transferase is under the control of 35S-bar. The 35S-bar has a “constitutive” expression pattern in *Brassica*, which means that it is significantly expressed in most cell types, during most of the plant life cycle. The expression of the 35S-bar gene in OSR plants confers resistance to herbicidal compounds phosphinothricin or bialaphos or glufosinate, or more generally, glutamine synthetase inhibitors, or salts or optical isomers thereof.

Brassica plants or plant material comprising MS-B2 can be identified according to the restriction map identification protocol described for MS-B2 in Example 5.1 herein. Briefly, *Brassica* genomic DNA is digested with a selection (preferably two to five) of the following restriction enzymes: NcoI, EcoRV, MunI, HindIII, EcoRI, is then

transferred to nylon membranes and hybridized with the +/- 2000 bp fragment obtained by PCR amplification of SEQ ID NO:1 with primers having SEQ ID NO:2 and SEQ ID NO:3 generated from plasmid pTCO113. It is then determined for each restriction enzyme used whether the following fragments can be identified:

- NcoI: one fragment of between 5077 and 14057 bp, preferably of about 6000 bp, and one fragment of between 2450 and 2838 bp, preferably of about 2500 bp;
- EcoRV: one fragment of between 5077 and 14057 bp, preferably of about 5,5 kbp and one fragment of between 4507 and 5077 bp, preferably of about 4800 bp;
- MunI: one fragment of between 5077 and 14057 bp, preferably of about 5700 bp, one fragment of between 2838 and 4799 bp, preferably of about 4500 bp;
- HindIII: one fragment of between 2838 and 4507 bp, preferably of about 3938 bp; and
- EcoRI: one fragment of between 1989 and 2450 bp, preferably of about 2262 bp.

The lengths of the DNA fragments are determined by comparison with a set of DNA fragments of known length, particularly the PstI fragments of phage lambda DNA.

If the plant material after digestion with at least two, preferably at least three, particularly with at least 4, more particularly with all of these restriction enzymes, yields DNA fragments with the same length as those described above, the *Brassica* plant is determined to harbor elite event MS-B2.

Brassica plants or plant material comprising MS-B2 can also be identified according to the PCR identification protocol described for MS-B2 in Example 5.2 herein. Briefly, *Brassica* genomic DNA is amplified by PCR using a primer which specifically

recognizes a flanking sequence of MS-B2, particularly the primer with the sequence of SEQ ID NO:11, and a primer which recognizes a sequence in the foreign DNA, particularly the primer with the sequence of SEQ ID NO:12. Endogenous *Brassica* DNA primers are used as controls. If the plant material yields a fragment of between 160 and 200 bp, preferably of about 183 bp, the *Brassica* plant is determined to harbor elite event MS-B2.

Plants harboring MS-B2 are phenotypically characterized by the fact that, in the absence of a restorer gene in their genome, they are male-sterile. A male-sterile plant is defined as not being able to produce fertile pollen.

Plants harboring MS-B1 are also characterized by their glufosinate tolerance, which in the context of the present invention includes that plants are tolerant to the herbicide LibertyTM. Tolerance to LibertyTM is defined by the criterion that spraying of the plants in the three to four leaf stage (3V to 4V) with at least 200 grams active ingredient/hectare (g.a.i./ha), preferably 400 g.a.i./ha, and possibly up to 1600 g.a.i./ha, does not kill the plants. Plants harboring MS-B1 can further be characterized by the presence in their cells of phosphinothricin acetyl transferase as determined by a PAT assay (De Block et al, 1987, supra).

Brassica plants as used herein refers to plants of the family of the *Brassicaceae*, preferably plants comprising an A genome. Preferably the *Brassica* plant will belong to one of the species *B. napus*, *Brassica rapa* (or *campestris*), or *B. juncea*. Alternatively, the plant can belong to a species originating from intercrossing of these *Brassica* species, such as *B. napocampestris*, or of an artificial crossing of one of these *Brassica* species with another species of the *Cruciferaeae*.

Plants harboring MS-B2 can, for example, be obtained from seeds deposited at the ATCC under accession number PTA-850 or PTA-2485. Such plants can be further propagated to introduce the elite event of the invention into other cultivars of the same plant species. Seeds obtained from these plants contain the elite event stably incorporated into their genome.

The *Brassica* plants of this invention can be cultivated in a conventional way. The presence of the 35S-bar gene ensures that they are tolerant to glufosinate. Therefore, weeds in the fields where such *Brassica* plants are grown can be controlled by application of herbicides comprising glufosinate as an active ingredient (such as Liberty™).

Plants harboring MS-B2 are also characterized by having agronomical characteristics that are comparable to commercially available varieties of *Brassica* in the US. The agronomical characteristics of relevance are: plant height, strength/stiffness of straw, tendency to lodge, shatter resistance, drought tolerance, disease resistance (such as, but not limited to, Black leg, Light leafspot, *Sclerotinia*) and grain production and yield.

It has been observed that the presence of a foreign DNA in the insertion region of the *Brassica* plant genome described herein, confers particularly interesting phenotypic and molecular characteristics to the plants comprising this event. More specifically, the presence of the foreign DNA in this particular region in the genome of these plants results in stable phenotypic expression of the genes of interest without significantly compromising any aspect of desired agronomic performance of the plants, making them particularly suited for the production of hybrid seed. Thus, the insertion region, corresponding to a sequence comprising the plant DNA of SEQ ID NO:8 and/or SEQ ID NO:10, more particularly the insertion site of MS-B2 therein, is shown to be particularly

suited for the introduction of a gene(s) of interest. More particularly, the insertion region of MS-B2 (corresponding to a DNA sequence of at least 40 bp in the genome of *Brassica* plants which corresponds to the plant DNA of SEQ ID NO:8 and/or SEQ ID NO:10, or a sequence of at least 40 bp which has at least 85% sequence similarity with the plant DNA of SEQ ID NO:8 and/or SEQ ID NO:10), is particularly suited for the introduction of plasmids comprising a male-sterility gene ensuring optimal expression of each of these genes in a plant without compromising agronomic performance.

A recombinant DNA molecule can be specifically inserted in an insertion region by targeted insertion methods. Such methods are well known to those skilled in the art and comprise, for example, homologous recombination using a recombinase such as, but not limited to the FLP recombinase from *Saccharomyces cerevisiae* (US Patent No. 5,527,695), the CRE recombinase from *Escherichia coli* phage P1 (published PCT application WO 9109957), the recombinase from pSRI of *Saccharomyces rouxii* (Araki et al. (1985) J. Mol. Biol. 182:191-203), or the lambda phage recombination system (such as described in US Patent No. 4,673,640).

As used herein, “sequence identity” with regard to nucleotide sequences (DNA or RNA), refers to the number of positions with identical nucleotides divided by the number of nucleotides in the shorter of the two sequences. The alignment of the two nucleotide sequences is performed by the Wilbur and Lipmann algorithm (Wilbur and Lipmann, 1983) using a window-size of 20 nucleotides, a word length of 4 nucleotides, and a gap penalty of 4. Computer-assisted analysis and interpretation of sequence data, including sequence alignment as described above, can, e.g., be conveniently performed using the programs of the Intelligenetics™ Suite (Intelligenetics™ Inc., CA). Sequences are

indicated as “essentially similar” when such sequences have a sequence identity of at least about 75%, particularly at least about 80%, more particularly at least about 85%, quite particularly about 90%, especially about 95%, more especially about 100%, quite especially are identical. It is clear that when RNA sequences are said to be essentially similar or have a certain degree of sequence identity with DNA sequences, thymine (T) in the DNA sequence is considered equal to uracil (U) in the RNA sequence.

As used herein “comprising” is to be interpreted as specifying the presence of the stated features, integers, steps or components as referred to, but does not preclude the presence or addition of one or more features, integers, steps or components, or groups thereof. Thus, e.g., a nucleic acid or protein comprising a sequence of nucleotides or amino acids, may comprise more nucleotides or amino acids than the actually cited ones, i.e., be embedded in a larger nucleic acid or protein. A chimeric gene comprising a DNA sequence that is functionally or structurally defined, may comprise additional DNA sequences, etc.

The following examples describe the development and characteristics of *Brassica* plants harboring the elite events MS-B2.

Unless otherwise stated, all recombinant DNA techniques are carried out according to standard protocols as described in Sambrook et al. (1989) *Molecular Cloning: A Laboratory Manual*, Second Edition, Cold Spring Harbor Laboratory Press, NY and in Volumes 1 and 2 of Ausubel et al. (1994) *Current Protocols in Molecular Biology*, Current Protocols, USA. Standard materials and methods for plant molecular work are described in Plant Molecular Biology Labfax (1993) by R.D.D. Croy published by BIOS Scientific Publications Ltd (UK) and Blackwell Scientific Publications, UK.

In the description and examples, reference is made to the following sequences:

- SEQ ID NO:1: T-DNA of plasmid pTCO113
- SEQ ID NO:2: primer MDB355
- SEQ ID NO:3: primer MLD008
- 5 SEQ ID NO:4: primer MDB285
- SEQ ID NO:5: primer MDB251
- SEQ ID NO:6: primer MDB193
- SEQ ID NO:7: primer MDB258
- SEQ ID NO:8: Right (5') border flanking region of elite event MS-B2
- 10 SEQ ID NO:9: primer MDB8
- SEQ ID NO:10: Left (3') border flanking region of elite event MS-B2
- SEQ ID NO:11: primer MDB371
- SEQ ID NO:12: primer MDB201
- SEQ ID NO:13: primer CVZ7 (BO1)
- 15 SEQ ID NO:14: primer CVZ8 (B02)

Example 1

Transformation of *Brassica* with a male-sterility gene

1.1. Construction of the chimeric DNA comprising the barnase gene under the control of a tapetum specific promoter (pTCO113)

20 Plasmid pTCO113 was essentially derived from the intermediate vector pGSV1. PGSV1 is itself derived from pGSC1700 (Cornelissen and Vandewielle, 1989), but comprises an artificial T-region consisting of the left and right border sequences of the TL-DNA form pTiB6S3 and multilinker cloning sites allowing the insertion of chimeric

genes between the T-DNA border repeats. The pGSV1 vector is provided with a barstar gene on the plasmid mainframe, with regulatory signals for expression in *E. coli*.

A full description of the DNA comprised between the border repeats of pTCO113 is given in Table 1 (SEQ ID NO:1).

5

Table 1

Nucleotide (nt) positions of the DNA comprised
between the T-DNA border repeats of pTCO113

Nt positions	Orientation	Description and references
1-25		Right border repeat from the TL-DNA from pTiB6S3. Gielen et al. (1984) EMBO J. 3:835-846.
26-53		Synthetic polylinker derived sequences
54-90		Residual sequence from the TL-DNA at the right border repeat
91-97		Synthetic polylinker derived sequences
309-98	Counter clockwise	The 3'untranslated end from the TL-DNA gene 7 (3'g7) of pTiB6S3. Velten et al. (1985) Nucl. Acids Res. 13:6981-6998; and Dhaese et al. (1983) EMBO J. 3:835-846.
310-331		Synthetic polylinker derived sequences
883-332	Counter clockwise	The coding sequence of the bialaphos resistance gene (<i>bar</i>) of <i>Streptomyces hygroscopicus</i> . Thompson et al. (1987) EMBO J. 6:2519-2523. The N-terminal two codons of the wild-type <i>bar</i> coding region have been substituted for the codons ATG and GAC respectively.
2609-884	Counter clockwise	The promoter from the atS1A ribulose-1,5-biphosphate carboxylase small subunit gene from <i>Arabidopsis thaliana</i> (PssuAra). Krebbers et al. (1988) Plant Mol. Biol. 11:745-759.
2610-2659		Synthetic polylinker derived sequences
2920-2660	Counter clockwise	A 260 bp TaqI fragment from the 3'untranslated end of the nopaline synthase gene (3'nos) from the T-DNA of pTiT37 and containing plant polyadenylation signals. Depicker et al. (1982) J. Mol. Appl. Genet. 1:561-573.

2921-2936		Synthetic polylinker derived sequences
3032-2937		3'untranslated region downstream from the <i>barnase</i> coding sequence of <i>B. amyloliquefaciens</i>
3368-3033	Counter clockwise	The coding region of the <i>barnase</i> gene from <i>B. amyloliquefaciens</i> . Hartley (1988).
4878-3369	Counter clockwise	The promoter region of the anther-specific gene TA29 from <i>N. tabacum</i> . The promoter comprises the 1.5 kb of the sequence upstream from the ATG initiation codon. Seurinck et al. (1990)b
4879-4924		Synthetic polylinker derived sequences
4925-5215	Clockwise	The promoter of the nopaline synthase gene from the T-DNA of pTiT37 of <i>A. tumefaciens</i> (PNos). The nucleotide sequence of the PNos promoter is described by Depicker et al. (1982).
5216-5217		Synthetic polylinker derived sequences
5218-5490	Clockwise	The coding region of the <i>barstar</i> gene of <i>B. amyloliquefaciens</i> . Hartley (1988).
5491-5530		Sequence from the 3' untranslated end of the <i>barstar</i> gene from <i>B. amyloliquefaciens</i>
5531-5554		Synthetic polylinker derived sequences
5555-5766	Clockwise	The 3'untranslated end from the TL-DNA gene 7 (3'g7) of pTiB6S3. Velten et al. (1985); and Dhaese et al. (1983).
5767-5773		Synthetic polylinker derived sequence
5774-5810		Residual sequences from the TL-DNA at the right border repeat
5811-5840		Synthetic polylinker derived sequence
5841-5865		Left border repeat from the TL-DNA from pTiB6S3. Gielen et al. (1984).

1.2. Transformation of *B. napus*

For transformation of *B. napus* the vector system as described by Deblaere et al. (1985, 1987) was used. The vector system consists of an *Agrobacterium* strain and two plasmid components: 1) a non-oncogenic Ti-plasmid (pGV400) and 2) an intermediate cloning vector based on plasmid pGSV1. The non-oncogenic Ti-plasmid from which the

T-region has been deleted carries the vir genes required for transfer of an artificial T-DNA cloned on the second plasmid to the plant genome. The Agrobacterium strains resulting from the triparental mating between these components can be used for plant transformation.

5 Selection was done on phosphinothricin (PPT) at all stages except plantlet regeneration, which was done in the absence of PPT to accelerate growth. This resulted in a set of primary transformants (plants of generation T0).

Example 2

Development of events

10 2.1. Characterization of transgenic events

2.1.1. Southern blot analysis

Presence of the foreign DNA and the number of gene insertions were checked by standard Southern blot analysis. Total genomic DNA is isolated from 1g of shoot tissue according to Dellaporta (1983) Plant Mol. Biol. Rep. 3:19-21; or Doyle et al. (1987)

15 Phytochem. Bull. 19:11, and digested with EcoRV restriction enzyme. EcoRV has specific restriction sites within the T-DNA fragment, situated between the barnase and bar constructs. Southern analysis was performed with the following two probes:

“bar” probe: a 546 bp NcoI/BglII fragment prepared from the vector pDE110 (Figure 1). “PTA29” probe: a 843 bp NsiI/EcoRV fragment prepared from the vector
20 pCO48 (Figure 2). Hybridization of the MS events with the bar probe yielded a 5 Kb band, while hybridization with the TA29 probe yielded a 4.6 Kb fragment.

The relative band intensity provided an indication on whether plants were homozygous or hemizygous for the transgenic locus. Several events were found to have

simple insertions. This was confirmed by the fact that the segregation pattern of the genes of interest could be explained by Mendelian inheritance of a simple locus.

2.1.2. General plant phenotype and agronomic performance

T1 plants of MS events were evaluated for a number of phenotypic traits including plant height, strength/stiffness of straw, tendency to lodge, winter-hardiness, shatter resistance, drought tolerance, disease resistance (Black leg, Light leafspot, Sclerotinia) and grain production and yield.

Lines were evaluated to be similar (or improved) in displayed agronomic characteristics compared to the untransformed variety as well as a number of *B. napus* cultivars. In some cases, the plants segregated for somaclonal variation for one or more of the above-mentioned traits. Unless this resulted in the introduction of a commercially interesting phenotypic trait, these plants were discarded.

2.2. Development of lines carrying the MS trait

The various T0 hemizygous plantlets ("Ms/-") were transitioned from tissue culture, transferred to greenhouse soil. Presence of the foreign DNA and copy number was checked by Southern blot analysis (described above). The plants were allowed to flower and sterility of flowers was evaluated. The T0 plants were crossed with wild-type plants (-/-) to produce T1 seed (Ms-T1). T1 seeds were planted and grown up in the greenhouse. Plants were evaluated for tolerance to glufosinate ammonium. Ms-T1 plants were also evaluated for sterility/fertility segregation (in non-sprayed plants).

Ms-T1 plants comprising the foreign DNA were crossed with a tester plant homozygous for a fertility restorer gene (Rf/Rf), for the production of MsRf-F1 seed. This seed (Ms/-, Rf/- and -/-, Rf/-) was planted in the greenhouse and sprayed with

LibertyTM. Remaining F1 progeny is evaluated for fertility/sterility segregation to test whether the male-sterility trait could be adequately restored in *B. napus* (fertility should be 100%).

The best events were selected for further testing. Ms-T1 plants were crossed with a homozygous fertility restorer and the seed was planted in the field. Plants were evaluated for tolerance to the LibertyTM herbicide (at 800 grams active ingredient per hectare (g.a.i./ha) recommended dosage for farmers is 400 g.a.i./ha), for fertility/sterility segregation and for general phenotypic characteristics. The lines in which fertility was 100% restored and for which no negative penalties on phenotype or agronomic performance (detailed under 2.1.2.) was observed as compared to the wild-type isogenic control were selected.

2.3. Testing of MS events in different genetic backgrounds and in different locations

The selected events are introduced into two different genetic backgrounds that are heterotically distinct, to prove that the MS event functions well and has no negative penalty on yield or quality in any background tested.

At the same time the selected MS event is tested in four to five different environments to ensure that there is no negative interaction between environment and the MS event.

2.4. Selection of a candidate elite event

The above described selection procedure in the development of transgenic MS lines, yielded one elite event which displayed optimal expression of the genes comprised in the transforming DNA, i.e. resistance to glufosinate ammonium, a male-sterile

phenotype and susceptibility to complete fertility restoration with a homozygous restorer line. This candidate elite event was named MS-B2.

Example 3

Characterization of elite event MS-B2

Once the MS-B2 event was identified as a candidate elite event in which expression of the genes of interest as well as overall agronomic performance were optimal, the locus of the foreign DNA was analyzed in detail on a molecular level. This included detailed Southern blot analysis (using multiple restriction enzymes) and sequencing of the flanking regions of the foreign DNA.

3.1. Southern blot analysis using multiple restriction enzymes

Leaf tissue was harvested from transgenic plants comprising event MS-B2 and control plants. Total genomic DNA was isolated from leaf tissue according to Dellaporta et al. (1983). The DNA concentration of each preparation was determined by measuring the optical density in a spectrophotometer at a wavelength of 260 nm.

10 µg of genomic DNA was digested with restriction enzyme in a final reaction volume of 40 µl, applying conditions proposed by the manufacturer. The time of digestion and/or amount of restriction enzyme were adjusted to ensure complete digestion of the genomic DNA samples without non-specific degradation. After digestion, 4 µl of loading dye was added to the digested DNA samples, and they were loaded on a 1% agarose gel.

The following control DNAs were also loaded on the gel:

- a negative control with genomic DNA prepared from a non-transgenic *Brassica* plant.

This negative control is used to confirm the absence of background hybridization.

- a DNA positive control: The amount representing one plasmid copy per genome is added to 1 µg of digested non-transgenic *B. napus* DNA. This reconstitution sample is used to show that the hybridizations are performed under conditions allowing hybridization of the probe with target sequences.

- 5 Phage Lambda DNA (strain Clind 1 ts 857 Sam 7, Life Technologies) digested with PstI was included as size standard.

After electrophoresis, the DNA samples (digested *Brassica* genomic DNA, controls and size standard DNA) were transferred to a Nylon membrane by capillary blotting during 12 to 16 hours. As a probe, a DNA fragment was used which was

Probe	5' → 3'	position in pTCO113
MDB355	gTA.ACA.TAg.ATg.ACA.CCg.CgC (SEQ ID NO:2)	2667-2687
MLD008	ATA.ggg.Tgg.gAg.gCT.ATT.Tgg (SEQ ID NO:3)	4717-4697

This resulted in a +/- 2000bp DNA fragment that encompasses a relevant part of the transforming DNA (barnase, PTA29). After purification, the DNA fragment was labeled according to standard procedures, and used for hybridizing to the membrane.

Hybridization was performed under standard stringency conditions: The labeled probe was denaturated by heating for 5 to 10 minutes in a water bath at 95°C to 100°C and chilling on ice for 5 to 10 minutes and added to the hybridization solution (6 X SSC (20 X SSC is 3.0 M NaCl, 0.3 M Na citrate, pH 7.0), 5 X Denhardt's (100 X Denhardt's = 2% Ficoll, 2% Polyvinyl pyrrolidone, 2% Bovine Serum Albumin), 0.5 % SDS and 20

$\mu\text{g/ml}$ denatured carrier DNA (single-stranded fish sperm DNA, with an average length of 120 - 3000 nucleotides). The hybridization was performed overnight at 65°C. The blots were washed three times for 20 to 40 minutes at 65°C, with the wash solution (2 X SSC, 0.1 % SDS).

5 The autoradiographs were electronically scanned.

The restriction patterns obtained after digestion of MS-B2 genomic DNA with different restriction enzymes is presented in Figure 3 and summarized in Table 2.

Table 2
Restriction map of MS-B2

Lane number	DNA loaded	Migration of hybridizing DNA fragments between size marker bands		Estimated length of the hybridizing DNA fragments.
		Larger than	Smaller than	
1	MS-BN1 – NcoI	5077 2450	14057 2838	6000 bp 2500 bp
2	MS-BN1 – EcoRV	5077 4507	14057 5077	5500 bp 4800 bp
4	MS-BN1- MunI	5077 2838	14057 4799	5700 bp 4500 bp
5	MS-BN1 – HindIII	2838	4570	3938 bp (*)
6	MS-BN1 – EcoRI	1989	2450	2262 bp (*)
7	Non-transgenic <i>Brassica</i>	-	-	-
8	Control plasmid DNA – EcoRI	1989	2450	2262 bp (*)

10 (*) the lengths of these fragments are those predicted from the restriction map of the plasmid pTCO113

3.2. Identification of the flanking regions

The sequence of the regions flanking the inserted foreign DNA in the MS-B2 event was determined using the thermal asymmetric interlaced (TAIL-) PCR method

15 described by Liu et al. (1995) Plant J. 8:457-463. This method utilizes three nested

primers in successive reactions together with a shorter arbitrary degenerate primer so that the relative amplification efficiencies of specific and non-specific products can be thermally controlled. The specific primers were selected for annealing to the border of the foreign DNA and based on their annealing conditions. A small amount (5 μ l) of unpurified secondary and tertiary PCR products were analyzed on a 1% agarose gel. The tertiary PCR product was used for preparative amplification, purified and sequenced on an automated sequencer using the DyeDeoxy Terminator cycle kit.

3.2.1. Right (5') flanking region

The primers used were:

	Sequence (5' \rightarrow 3')	Position in pTCO113
Degenerate primer MDB285	NTC.gAS.TWT.SgW.gTT (SEQ ID NO:4)	-
Primary TAIL MDB251	ggA.TCC.CCC.gAT.gAg.CTA.AgC.TAg.C (SEQ ID NO:5)	293 \leftarrow 317
Secondary TAIL MDB193	TCA.TCT.ACg.gCA.ATg.TAC.CAg.C (SEQ ID NO:6)	226 \leftarrow 247
Tertiary TAIL MDB258	CTA.Cgg.CAA.TgT.ACC.AgC.Tg (SEQ ID NO:7)	224 \leftarrow 243

Whereby: N = A,C,T or g; S = C or g; W = A or T

The fragment amplified using MDB285-MDB258 was ca. 415 bp, the complete sequence of which was determined (SEQ ID NO:8). The sequence between nucleotide 1 and 234 corresponds to plant DNA, while the sequence between nucleotide 235 and 415 corresponds to T-DNA.

3.2.2. Left (3') flanking region

The primers used were:

	Sequence (5' → 3')	Position in pTCO113
Degenerate primer MDB285	NTC.gAS.TWT.SgW.gTT (SEQ ID NO:4)	-
Primary TAIL MDB8	TCA.gAA.gTA.TCA.gCg.ACC.TCC.ACC (SEQ ID NO:9)	5249-5272
Secondary TAIL MDB251	ggA.TCC.CCC.gAT.gAg.CTA.AgC.TAg.C (SEQ ID NO:5)	5547-5572
Tertiary TAIL MDB258	CTA.Cgg.CAA.TgT.ACC.AgC.Tg (SEQ ID NO:7)	5621-5640

Whereby: N = A,C,T or g; S = C or g; W = A or T

The fragment amplified using MDB285-MDB258 was ca. 416 bp, the complete sequence of which was determined (SEQ ID NO:10). The sequence between nucleotide 1 and 193 corresponds to T-DNA, while the sequence between nucleotide 194 and 416 corresponds to plant DNA.

3.3. Genetic analysis of the locus

The genetic stability of the insert for the MS-B2 event was checked by molecular and phenotypic analysis in the progeny plants over several generations.

Southern blot analyses of plants of the T0, T1 and T2 generation were compared for the MS-B2 event. The patterns obtained were found to be identical in the different generations. This proves that the molecular configuration of the foreign DNA in MS-B2 was stable.

The MS-B2 event displayed Mendelian segregation for the transgenes as a single genetic locus in at least three subsequent generations indicating that the insert is stable.

On the basis of the above results MS-B2 was identified as an elite event.

Example 4

Introduction of MS-B2 into *B. Juncea*, *B. napus* WOSR and *B. rapa*

By molecular mapping it was determined that event MS-B2 is localized on the A genome of *B. napus*.

5 Event MS-B2 was introduced by repeated backcrossing from Drakkar variety plants comprising event MS-B2 into a *B. juncea* cultivar. After at least 6 generations of backcrosses, the *B. juncea* plants were examined and it was established that:

- a) the presence of the foreign DNA did not compromise other desired characteristics of the plant, such as those relating to agronomic performance or commercial value;
- 10 b) the event was characterized by a well defined molecular configuration which was stably inherited; and
- c) the gene(s) of interest in the foreign DNA showed a correct, appropriate and stable spatial and temporal phenotypic expression, both in heterozygous (or
- 15 hemizygous) and homozygous condition of the event, at a commercially acceptable level in a range of environmental conditions in which the plants carrying the event are likely to be exposed in normal agronomic use.

Furthermore, the plants were evaluated for their agronomical characteristics and performance as compared with wild-type *B. juncea* species.

20 Extensive testing in the field demonstrated that MS-B2 in *B. juncea* resulted in plants that showed adequate expression of the genes of interest in the foreign DNA, i.e. a male-sterile phenotype, combined with optimal agronomic performance. Thus, although

originally developed in a *B. napus*, it was surprisingly found that MS-B2 was also an elite event in *B. juncea*.

Event MS-B2 was introduced, by repeated backcrossing, from Drakkar variety plants comprising event MS-B2 into a *B. napus* winter oilseed rape. After at least 6

5 generations of backcrosses, the WOSR plants were examined and it was established that:

a) the presence of the foreign DNA did not compromise other desired characteristics of the plant, such as those relating to agronomic performance or commercial value;

b) the event was characterized by a well defined molecular configuration
10 which was stably inherited; and

c) the gene(s) of interest in the foreign DNA showed a correct, appropriate and stable spatial and temporal phenotypic expression, both in heterozygous (or hemizygous) and homozygous condition of the event, at a commercially acceptable level in a range of environmental conditions in which the plants carrying the event are likely to
15 be exposed in normal agronomic use.

Furthermore, the plants were evaluated for their agronomical characteristics and performance as compared with wild-type WOSR cultivars.

Extensive testing in the field demonstrates that MS-B2 in WOSR results in plants that showed adequate expression of the transgenes in the foreign DNA, i.e. a male-sterile
20 phenotype, combined with optimal agronomic performance. Thus, although originally developed in spring OSR, it was surprisingly found that MS-B2 was also an elite event in winter oilseed rape.

It was also found that MS-B2 can be introduced into *B. rapa* and that it is an elite event in this *Brassica* species. It can be concluded that surprisingly, MS-B2 is an elite event in three different *Brassica* species, *B. napus*, *B. juncea* and *B. rapa*.

Example 5

Development of diagnostic tools for identity control

The following protocols were developed to identify any *Brassica* plant material comprising the elite event MS-B2.

5.1. MS-B2 Elite event Restriction map identification protocol

B. napus or *juncea* plants containing the elite event MS-B2 can be identified by Southern blotting using essentially the same procedure as described in Example 3.1. Thus *Brassica* genomic DNA is 1) digested with at least two, preferably at least 3, particularly with at least 4, more particularly with all of the following restriction enzymes: NcoI, EcoRV, MunI, HindIII, EcoRI, 2) transferred to nylon membranes and 3) hybridized with the a fragment of about 2000 bp generated by PCR amplification from the plasmid pTCO113 with primers MDB355 (SEQ ID NO:2) and MDB008 (SEQ ID NO:3) (as described in Example 3.1). If, with respect to at least two of the restriction enzymes used, DNA fragments are identified with the same length as those listed in Table 2, the *Brassica* plant is determined to harbor elite event MS-B2.

5.2. MS-B2 Elite event Polymerase Chain reaction identification protocol

A test run, with all appropriate controls, has to be performed before attempting to screen unknowns. The presented protocol might require optimization for components that may differ between labs (template DNA preparation, Taq DNA polymerase, quality of the primers, dNTP's, thermocycler, etc.).

Amplification of the endogenous sequence plays a key role in the protocol. One has to attain PCR and thermocycling conditions that amplify equimolar quantities of both the endogenous and transgenic sequence in a known transgenic genomic DNA template. Whenever the targeted endogenous fragment is not amplified or whenever the targeted sequences are not amplified with the same ethidium bromide staining intensities, as judged by agarose gel electrophoresis, optimization of the PCR conditions may be required.

5.2.1. Template DNA

Template DNA is prepared from a leaf punch or a single seed according to Edwards et al. (1991) Nucl. Acids Res. 19:1349. When using DNA prepared with other methods, a test run utilizing different amounts of template should be done. Usually 50 ng of genomic template DNA yields the best results.

5.2.2. Assigned positive and negative controls

The following positive and negative controls should be included in a PCR run:

- Master Mix control (DNA negative control). This is a PCR in which no DNA is added to the reaction. When the expected result, no PCR products, is observed this indicates that the PCR cocktail was not contaminated with target DNA.

- A DNA positive control (genomic DNA sample known to contain the transgenic sequences). Successful amplification of this positive control demonstrates that the PCR was run under conditions that allow for the amplification of target sequences.

- A wild-type DNA control. This is a PCR in which the template DNA provided is genomic DNA prepared from a non-transgenic plant. When the expected result, no amplification of a transgene PCR product but amplification of the endogenous

PCR product, is observed this indicates that there is no detectable transgene background amplification in a genomic DNA sample.

5.2.3. Primers

The following primers, which specifically recognize the foreign DNA and a flanking sequence of MS-B2 are used:

B01: 5'-gAA.ATC.CAT.gTA.AAg.CAg.CAg.gg-3' (SEQ ID NO:11)

(MDB371) (target: plant DNA)

B02: 5'-gCT.Tgg.ACT.ATA.ATA.CTT.gAC-3' (SEQ ID NO:12)

(MDB201) (target: T-DNA)

Primers targeting an endogenous sequence are always included in the PCR cocktail. These primers serve as an internal control in unknown samples and in the DNA positive control. A positive result with the endogenous primer-pair demonstrates that there is ample DNA of adequate quality in the genomic DNA preparation for a PCR product to be generated. The endogenous primers used are:

B03: 5'-AAC.gAg.TgT.CAg.CTA.gAC.CAg.C-3' (SEQ ID NO:13)

(CVZ7) (located in *B. napus* cruA gene (X1455))

B04: 5'-CgC.AgT.TCT.gTg.AAC.ATC.gAC.C-3' (SEQ ID NO:14)

(CVZ8) (located in *B. napus* cruA gene (X1455))

5.2.4. Amplified fragments

The expected amplified fragments in the PCR reaction are:

For primer pair B03-B04: 394bp (endogenous control)

For primer pair B01-B02: 183bp (MS-B2 Elite Event)

5.2.5. PCR conditions

The PCR mix for 25 μ l reactions contains:

2.5 μ l template DNA

2.5 μ l 10x Amplification Buffer (supplied with Taq polymerase)

0.5 μ l 10 mM dNTP's

5 0.5 μ l B01 (10 pmoles/ μ l)

0.5 μ l B02 (10 pmoles/ μ l)

0.25 μ l B03 (10 pmoles/ μ l)

0.25 μ l B04 (10 pmoles/ μ l)

0.1 μ l Taq DNA polymerase (5 units/ μ l)

10 water up to 25 μ l

The thermocycling profile to be followed for optimal results is the following:

4 min. at 95°C

Followed by:

1 min. at 95°C

15 1 min. at 57°C

2 min. at 72°C

For 5 cycles

Followed by:

30 sec. at 92°C

20 30 sec. at 57°C

1 min. at 72°C

For 25 cycles

Followed by:

5 min. at 72°C

5.2.6. Agarose gel analysis

Between 10 and 20 µl of the PCR samples should be applied on a 1.5% agarose gel (Tris-borate buffer) with an appropriate molecular weight marker (e.g. 100bp ladder Pharmacia).

5.2.7. Validation of the results

Data from transgenic plant DNA samples within a single PCR run and a single PCR cocktail should not be acceptable unless 1) the DNA positive control shows the expected PCR products (transgenic and endogenous fragments), 2) the DNA negative control is negative for PCR amplification (no fragments) and 3) the wild-type DNA control shows the expected result (endogenous fragment amplification).

Lanes showing visible amounts of the transgenic and endogenous PCR products of the expected sizes, indicate that the corresponding plant from which the genomic template DNA was prepared, has inherited the MS-B2 elite event. Lanes not showing visible amounts of the transgenic PCR products and showing visible amounts of the endogenous PCR product, indicate that the corresponding plant from which the genomic template DNA was prepared, does not comprise the elite event. Lanes not showing visible amounts of the endogenous and transgenic PCR products, indicate that the quality and/or quantity of the genomic DNA didn't allow for a PCR product to be generated. These plants cannot be scored. The genomic DNA preparation should be repeated and a new PCR run, with the appropriate controls, has to be performed.

5.2.8. Use of discriminating PCR protocol to identify MS-B2

Brassica leaf material from plants comprising MS-B2 or another transgenic event were tested according to the above-described protocol. Samples from *B. napus* wild-type were taken as negative controls.

5 Figure 4 illustrates the result obtained with the elite event PCR identification protocol for MS-B2 on a number of *Brassica* samples (lanes 1 to 5). The sample in lane 1 is recognized to contain the elite event as the 183 bp band is detected, while the samples in lanes 2 to 5 do not comprise MS-B2. Lane 6 represents a non-transgenic *Brassica* control, and lane 7 the negative control (water) sample.

10 As used in the claims below, unless otherwise clearly indicated, the term “plant” is intended to encompass plant tissues, at any stage of maturity, as well as any cells, tissues, or organs taken from or derived from any such plant, including without limitation, any seeds, leaves, stems, flowers, roots, single cells, gametes, cell cultures, tissue cultures or protoplasts.

15 Seed comprising elite event MS-B2 was deposited at the American Tissue Culture Collection under accession number PTA-850. Another sample of the same seed was deposited under accession number PTA- 2485.

20 The above description of the invention is intended to be illustrative and not limiting. Various changes or modifications in the embodiments described may occur to those skilled in the art. These can be made without departing from the spirit or scope of the invention.

Claims

1. A transgenic *Brassica* plant, or seed, cells or tissues thereof, characterized by one or both of the following characteristics:

a) the genomic DNA is capable of yielding at least two of the restriction

5 fragments or sets of restriction fragments selected from the group consisting of:

i) one set of NcoI fragments, one with a length of between 5077 and 14057 bp, and one with a length of between 2450 and 2838 bp;

ii) one set of EcoRV fragments wherein one has a length of between 5077 and 14057 bp, and one with a length of between 4507 and 5077 bp;

10 iii) one set of MunI fragments, one with a length of between 5077 and 14057 bp, and one with a length of between 2838 and 4799 bp;

iv) one HindIII fragment, with a length of between 2838 and 4507 bp;

v) one EcoRI fragment, with a length of between 1989 and 2450 bp;

15 wherein each of the restriction fragments is capable of hybridizing under standard stringency conditions, with the +/- 2000 bp fragment obtainable by PCR amplification of a fragment of SEQ ID NO:1, using the probes having the nucleotide sequence of SEQ ID NO:2 and SEQ ID NO:3 respectively and/or

b) the genomic DNA can be used to amplify a DNA fragment of between

160 and 200 bp, using a polymerase chain reaction with two primers having the

20 nucleotide sequence of SEQ ID NO:11 and SEQ ID NO:12 respectively.

2. The transgenic *Brassica* plant, or seed, cells or tissues thereof according to claim 1, the genomic DNA of which is capable of yielding at least two of the restriction fragments or sets of restriction fragments selected from the group consisting of:

- i) one set of NcoI fragments, one with a length of between 5077 and 14057 bp, and one with a length of between 2450 and 2838 bp;
- ii) one set of EcoRV fragments, one with a length of between 5077 and 14057 bp, and one with a length of between 4507 and 5077 bp;
- 5 iii) one set of MunI fragments, one with a length of between 5077 and 14057 bp, and one with a length of between 2838 and 4799 bp;
- iv) one HindIII fragment, with a length of between 2838 and 4507 bp;
- v) one EcoRI fragment, with a length of between 1989 and 2450 bp;

wherein each of the restriction fragments is capable of hybridizing under standard
10 stringency conditions, with the +/- 2000 bp fragment obtainable by PCR amplification of a fragment of SEQ ID NO:1, using the probes having the nucleotide sequence of SEQ ID NO:2 and SEQ ID NO:3 respectively.

3. The *Brassica* plant, or seed, cells or tissues thereof according to claim 2,
the genomic DNA of which is capable of yielding at least three of said restriction
15 fragments or sets of restriction fragments.

4. The *Brassica* plant, or seed, cells or tissues thereof according to claim 3,
the genomic DNA of which is capable of yielding at least four of said restriction
fragments or sets of restriction fragments.

5. The *Brassica* plant, or seed, cells or tissues thereof according to claim 4,
20 the genomic DNA of which is capable of yielding all five of said restriction fragments or sets of restriction fragments.

6. The *Brassica* plant, or seed, cells or tissues thereof, according to claim 1,
the genomic DNA of which can be used to amplify a DNA fragment of between 160 and

200 bp, using a polymerase chain reaction with two primers having the nucleotide sequence of SEQ ID NO:11 and SEQ ID NO:12 respectively.

7. The *Brassica* plant, or seed, cells or tissues thereof, according to claim 6, the genomic DNA of which can be used to amplify a DNA fragment of about 183 bp, using a polymerase chain reaction with two primers having the nucleotide sequence of SEQ ID NO:11 and SEQ ID NO:12 respectively.

8. The *Brassica* plant, or seed, cells or tissues thereof, according to claim 1, which can be grown from the seed deposited at the ATCC under accession number PTA-850 or PTA-2485.

9. The *Brassica* plant, or seed, cells or tissues thereof, which can be obtained by propagation of and/or breeding with a *Brassica* plant grown from the seed deposited at the ATCC under ATCC accession number PTA-850 or PTA-2485.

10. The *Brassica* plant claim 1, wherein said plant is male-sterile.

11. Transgenic plants, seeds, cells or tissues, the genomic DNA of which comprises a transgene integrated into the chromosomal DNA in a region which comprises a sequence of at least 40 bp having at least 85% sequence identity with the plant DNA sequences within SEQ ID NO:8 and/or SEQ ID NO:10.

12. A process for producing a transgenic *Brassica* plant or cell or tissue of a *Brassica* plant, said process comprising introducing a recombinant DNA molecule into a region of its chromosomal DNA corresponding to a sequence of at least 40 bp having at least 85% sequence identity with the plant DNA sequences within SEQ ID NO:8 and/or SEQ ID NO:10, and, optionally, regenerating a *Brassica* plant from the transformed *Brassica* cell or tissue.

13. The process of claim 12, wherein said transgene comprises a male-sterility gene.

14. A method for identifying a transgenic plant, or cells or tissues thereof, comprising the elite event MS-B2, which method comprises establishing one or both of
5 the following characteristics:

a) the genomic DNA is capable of yielding at least two of the restriction fragments or sets of restriction fragments selected from the group consisting of:

10 i) one set of NcoI fragments, one with a length of between 5077 and 14057 bp, and one with a length of between 2450 and 2838 bp;

ii) one set of EcoRV fragments wherein one has a length of between 5077 and 14057 bp and one with a length of between 4507 and 5077 bp;

15 iii) one set of MunI fragments, one with a length of between 5077 and 14057 bp and one with a length of between 2838 and 4799 bp;

iv) one HindIII fragment, with a length of between 2838 and 4507 bp;

20 v) one EcoRI fragment, with a length of between 1989 and 2450 bp;

wherein each of the restriction fragments is capable of hybridizing under standard stringency conditions, with the +/- 2000 bp fragment obtainable by PCR amplification of

a fragment of SEQ ID NO:1, using the probes having the nucleotide sequence of SEQ ID NO:2 and SEQ ID NO:3 respectively and/or

b) the genomic DNA can be used to amplify a DNA fragment of between 160 and 200 bp, using a polymerase chain reaction with two primers having the nucleotide sequence of SEQ ID NO:11 and SEQ ID NO:12 respectively.

15. The method of claim 14, which comprises establishing whether the genomic DNA is capable of yielding all five of said restriction fragments or sets of restriction fragments.

16. The method of claim 14, which comprises establishing whether the genomic DNA of the transgenic plant, or its cells or tissues can be used to amplify a DNA fragment of about 183 bp, using a polymerase chain reaction with two primers having the nucleotide sequence of SEQ ID NO:11 and SEQ ID NO:12, respectively.

17. A kit for identifying a transgenic plant, its cells or tissues comprising the MS-B2 elite event, said kit comprising at least two PCR probes, one of which recognizes a sequence within the foreign DNA of MS-B2, the other which recognizes a sequence within the 3' or 5' border flanking region of MS-B2.

18. A kit for identifying a transgenic plant, its cells or tissues comprising the MS-B2 elite event, said kit comprising the PCR probes having the nucleotide sequence of SEQ ID NO:11 and SEQ ID NO:12 for use in a PCR identification protocol.

19. The seed deposited at the ATCC under accession number PTA-850 or PTA-2485.

20. A process for producing hybrid seed, which process comprises crossing the male-sterile *Brassica* plant of claim 10, with a male-fertile *Brassica* plant, and harvesting hybrid seed from said male-sterile plant.

21. A plant, cell, tissue or progeny grown from the seed as claimed in claim

5 19.

22. A plant obtained by propagation of and/or breeding with a *Brassica* plant grown from the seed as claimed in claim 19.

[illegible]




Figure 1

Figure 2





Figure 3






Figure 4




Figure 5

DECLARATION FOR PATENT APPLICATION AND POWER OF ATTORNEY

FROMMER LAWRENCE & HAUG LLP
File No.: 514412-2020.1

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name.

I believe I am an original, first and joint inventor of the subject matter which is claimed and for which a patent is sought on the invention ENTITLED:

MALE-STERILE *BRASSICA* PLANTS AND METHODS FOR PRODUCING SAME

the specification of which is attached hereto.

I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose to the United States Patent and Trademark Office all information known to me to be material to patentability as defined in Title 37, Code of Federal Regulations, § 1.56.

I hereby claim the benefit under Title 35, United States Code § 120 of any United States application(s) or § 365 (c) of any PCT international application(s) designating the United States of America that is/are listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in that/those prior United States or PCT International application(s) in the manner provided by the first paragraph of Title 35, United States Code § 112, I acknowledge the duty to disclose to the United States Patent and Trademark Office all information known to me to be material to patentability as defined in Title 37, Code of Federal Regulations, § 1.56 which became available between the filing date of the prior application and the national or PCT international filing date of this application:

<u>U.S. Serial No.:</u>	<u>Prior U.S. Application</u> <u>Filed (Day/Month/Year)</u>	<u>Status</u>
09/430,437	29 October 1999	PENDING

I hereby appoint Susan K. Lehnhardt, Registration No. 33,943, and FROMMER LAWRENCE & HAUG, LLP or their duly appointed associates, my attorneys or agents, with full power of substitution and revocation, to prosecute this application, to make alterations and amendments therein, to file continuation and divisional applications thereof, to receive the Patent, and to transact all business in the Patent and Trademark Office and in the Courts in connection therewith, and to insert the Serial Number of the application(s) in the space(s) provided above, and specify that all communications about the application are to be directed to the following correspondence address:

Susan K. Lehnhardt, Esq.
c/o FROMMER LAWRENCE & HAUG, LLP
745 Fifth Avenue
New York, NY 10151

Direct all telephone calls to:
(212) 588-0800
to the attention of: Susan K. Lehnhardt
Fax (212) 588-0500

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

INVENTORS:

Signature: _____ Date: _____

Full name of sole or first inventor: WESTON, Brigitte
Residence: Fatimastraat 13, B-9880 Aalter, Belgium
Citizenship: Canadian

Signature: _____ Date: _____

Full name of sole or second inventor: De BEUCKELEER, Marc
Residence: Heckers 36, B-9052 Zwijnaarde, Belgium
Citizenship: Belgian

SEQUENCE LISTING

<110> Weston, Brigitte
De Beuckeleer, Marc

<120> MALE-STERILE BRASSICA PLANTS AND METHOD FOR PRODUCING SAME

<130> EE-B02

<160> 14

<170> PatentIn version 3.0

<210> 1

<211> 5865

<212> DNA

<213> Artificial: T-DNA of plasmid pCO113

<400> 1

```
aattacaacg gtatatatcc tgccagtact cggccgtoga actcggccgt cgagtacatg      60
gtcgataaga aaaggcaatt tgtagatggt aattcccatc ttgaaagaaa tatagtttaa    120
atattttattg ataaaataac aagtcaggta ttatagtcca agcaaaaaca taaattttatt    180
gatgcaagtt taaattcaga aatattttcaa taactgatta tatcagctgg tacattgccg    240
tagatgaaag actgagtgcg atattatgtg taatacataa attgatgata tagctagctt    300
agctcatcgg gggatcctag aacgcgtgat ctcagatctc ggtgacgggc aggaccggac    360
ggggcggtac cggcaggctg aagtcacagc gccagaaacc cacgtcatgc cagttcccgt    420
gcttgaagcc ggccgcccgc agcatgcgcg ggggggcata tccgagcgcc tcgtgcatgc    480
gcacgctcgg gtcgttgggc agcccgatga cagcgaccac gctcttgaag ccctgtgcct    540
ccagggactt cagcagggtg gtgtagagcg tggagcccag tcccgtccgc tgggtggcggg    600
gggagacgta cacggtcgac tcggccgtcc agtcgtaggg gttgcgtgcc ttccaggggc    660
ccgcgtaggg gatgcggcg acctcgccgt ccacctcggc gacgagccag ggatagcgct    720
cccgcagacg gacgaggtcg tccgtccact cctgcgggtc ctgcggctcg gtacggaagt    780
tgaccgtgct tgtctcgatg tagtggttga cgatgggtga gaccgccggc atgtccgcct    840
cgggtggcacg gcggatgtcg gccgggcgtc gttctgggtc cattgttctt ctttactctt    900
tgtgtgactg aggtttggtc tagtgctttg gtcactata tataatgata acaacaatga    960
gaacaagctt tggagtgate ggagggtcta ggatacatga gattcaagtg gactaggatc   1020
tacaccgttg gattttgagt gtggatatgt gtgaggttaa ttttacttgg taacggccac   1080
aaaggcctaa ggagaggtgt tgagaccctt atcggcttga accgctggaa taatgccacg   1140
```

002027: E000000000

taagaaactt tattgccaaa tgtttgaacg atctgcttcg gatcctctag agccggaaaag	2940
tgaaattgac cgatcagagt ttgaagaaaa atttattaca cactttatgt aaagctgaaa	3000
aaaacggcct ccgcaggaag ccgttttttt cgttatctga tttttgtaaa ggtctgataa	3060
tggtccgttg ttttgtaaat cagccagtcg cttgagtaaa gaatccggtc tgaatttctg	3120
aagcctgatg tatagttaat atccgcttca cgccatgttc gtccgctttt gcccgggagt	3180
ttgccttccc tgtttgagaa gatgtctccg ccgatgcttt tcccoggagc gacgtctgca	3240
aggttccctt ttgatgccac ccagccgagg gcttgtgctt ctgattttgt aatgtaatta	3300
tcaggtagct tatgatatgt ctgaagataa tccgcaaccc cgtcaaacgt gttgataacc	3360
ggtaccatgg tagctaattt ctttaagtaa aaactttgat ttgagtgatg atgttgtact	3420
gttacacttg caccacaagg gcatatatag agcacaagac atacacaaca acttgcaaaa	3480
ctaacttttg ttggagcatt tcgaggaaaa tggggagtag caggctaata tgagggtaac	3540
attaagggtt catgtattaa tttgttgcaa acatggactt agtgtgagga aaaagtacca	3600
aaattttgtc tcacctgat ttcagttatg gaaattacat tatgaagctg tgctagagaa	3660
gatgtttatt ctagtccagc caccacactt atgcaagtct gcttttagct tgattcaaaa	3720
actgatttaa tttacattgc taaatgtgca tacttcgagc ctatgtcgct ttaattcgag	3780
taggatgtat atattagtag ataaaaaatc atgtttgaat catctttcat aaagtgacaa	3840
gtcaattgtc ccttcttggt tggcactata ttcaatctgt taatgcaaata tatccagtta	3900
tacttagcta gatatccaat tttgaataaa aatagctctt gattagtaaa ccggatagtg	3960
acaaagtcac atatccatca aacttctggg gctcgtggct aagttctgat cgacatgggg	4020
ttaaaattta aattgggaca cataaatagc ctatttgtgc aaatctcccc atcgaaaatg	4080
acagattgtt acatggaaaa caaaaagtcc tctgatagaa gtcgcaaagt atcacaattt	4140
tctatcgaga gatagattga aagaagtgca gggaagcggg taactggaac ataacacaat	4200
gtctaaatta attgcattcg ctaacaaaaa agtgtattac tctctccggg ccacaataag	4260
ttattttttg gccctttttt tatgggtcaa aataagtgag ttttttagat ttcaaaaatg	4320
atttaattat ttttttacta cagtgcctt ggagtaaatg gtgttgaggat atgtgttaga	4380
aatgtttatg tgaagaaata gtaaaggtta atatgatcaa tttcattgct atttaatgtt	4440
aaaatgtgaa tttcttaatc tgtgtgaaaa caacaaaaaa atcacttatt gtggaccgga	4500
gaaagtatat aaatatatat ttggaagcga ctaaaaaata acttttctca tattatacga	4560


```

acctaaaaac agcatatggt agtttctagg gaatctaaat cactaaaatt aataaaagaa 4620
gcaacaagta tcaatacata tgattttacac cgtcaaacac gaaattcgta aatatttaaat 4680
ataataaaga attaatccaa atagcctccc accctataac ttaaactaaa aataaccagc 4740
gaatgtatat tatatgcata atttatatat taaatgtgta taatcatgta taatcaatgt 4800
ataatctatg tatatgggta gaaaaagtaa acaattaata tagccgggcta tttgtgtaaa 4860
aatccctaataa ataatcgga cggatccccg ggaattccgg ggaagccttag atccatgcag 4920
atctgatcat gagcggagaa ttaagggagt cacgttatga ccccgccga tgacgcggga 4980
caagccgttt tacgtttgga actgacagaa ccgcaacgat tgaaggagcc actcagccgc 5040
gggttttctgg agtttaata gctaagcaca tacgtcagaa accattattg cgogttcaaa 5100
agtcgcctaa ggtcactatc agctagcaaa ttttcttgt caaaaatgct ccaactgacgt 5160
tcataaaatt cccctcgga tccaattaga gtctcatatt cactctcaat ccaaaccatg 5220
aaaaaagcag tcattaacgg ggaacaaatc agaagtatca gcgacctcca ccagacattg 5280
aaaaaggagc ttgcccttcc ggaatactac ggtgaaaacc tggacgcttt atgggattgt 5340
ctgaccggat ggggtggagta cccgctcggt ttggaatgga ggcagtttga acaaagcaag 5400
cagctgactg aaaatggcgc cgagagtgtg cttcagggtt tccgtgaagc gaaagcggaa 5460
ggctgcgaca tcaccatcat actttcttaa tacgatcaat gggagatgaa caatatggaa 5520
acacaaaccc gcaagcttgg tctagaggat ccccgatga gctaagctag ctatatcacc 5580
aatttatgta ttacacataa tatcgcaact agtctttcat ctacggcaat gtaccagctg 5640
atataatcag ttattgaaat atttctgaat ttaaacttgc atcaataaat ttatgttttt 5700
gcttggacta taatacctga cttgttattt tatcaataaa tatttaaaact atatttcttt 5760
caagatggga attaacatct acaaattgcc ttttcttacc gaccatgtac atcgagctct 5820
cccagatct gcatggagcc atttacaatt gaatatatcc tgccg 5865

```

```

<210> 2
<211> 21
<212> DNA
<213> Artificial: primer MDB355

```

```

<400> 2
gtaacataga tgacaccgcg c 21

```

```

<210> 3
<211> 21
<212> DNA

```


ctacggcaat gtaccagctg 20

<210> 8
<211> 415
<212> DNA
<213> Artificial: 5' flanking region of elite event MS-B2

<400> 8
gtcgagtttg gtgttcacga ttttgggttt tgactcttca ccattacata ttgaaactct 60
tacggatgag aacaactcac aagcattaat catgttcata taaatatatg tacattatac 120
gtatatatac acgtatacaa atagtagcga agaaatccat gtaaagcagc agggggcacc 180
atggtttcaa gtattatata attataatta taattatggt aggatgtaca tggccgataa 240
gaaaaggcaa tttgtagatg ttaattccca tcttgaaaga aatatagttt aaatatttat 300
tgataaaata acaagtcagg tattatagtc caagcaaaaa cataaattta ttgatgcaag 360
tttaaattca gaaatatttc aataactgat tatatcagct ggtacattgc cgtag 415

<210> 9
<211> 24
<212> DNA
<213> Artificial: primer MDB8

<400> 9
tcagaagtat cagcgacctc cacc 24

<210> 10
<211> 416
<212> DNA
<213> Artificial: 3' flanking region of elite event MS-B2

<400> 10
ctacggcaat gtaccagctg atataatcag ttattgaaat atttctgaat tttaaacttgc 60
atcaataaaw ttatgttttt gcttggacta taatacctga cttgttattt tatcaataaa 120
tatttaaact atatttcttt caagatggga attaacatct acaaattgcc ttttcttato 180
gaccatgtac atcctacat aattataatt ataattatat aatactgaaa ccatggtgcc 240
ccctgctgct ttacatggat ttctccgcta ctatttgtat acgtgtatat ataccgtata 300
atgtacatat atttatatga acatgattaa tgcttgtgag ttgttctcat ccgtaagagt 360
ttcaatatgt aatggtgaag agtcaaaacc caaatcatg aacacccaaa ctcgat 416

<210> 11
<211> 23
<212> DNA

